

STUDY REPORT

1. Title: Acute Bacterial Meningitis in Thai Children

Principal Investigators: Chiraphun Duangmani, M.D.
Howard E. Noyes, Ph.D.
Sombat Singhurat, M.D.
Pramuan Sunakorn, M.D.

Associate Investigators: Udom Lexomboon, M.D.
Pethai Mansuvan, M.D.
Pitsawas Thongphakdi, M.D.

Assistant Investigators: Yachai Ampaipast, B.S.
Panyasri Benjadol, M.S.
Chantana Ratanavaraha, B.S.
Yongyuth Raengpradub, B.S.

OBJECTIVE

This is a continuation of a study of bacterial meningitis in children of Children's Hospital, Bangkok, Thailand. The objectives of this study are to determine the etiology of these infections, determine the antibiotic sensitivities of the causal agents and compare 3 antimicrobial regimens for the treatment of these patients.

DESCRIPTION

All patients with suspected bacterial meningitis admitted to Children's Hospital were included in this study. Antimicrobial therapy was based on order of admission. Daily administration consisted of (1) 10 megaunits of intravenous penicillin G sodium plus 50 mg/kg of intramuscular kanamycin sulfate or (2) the same dosage of penicillin plus 100 mg/kg of intramuscular chloramphenicol or (3) 150 mg/kg of intravenous sodium ampicillin. All regimens were divided into 4 doses which were given at 6 hour intervals.

Diagnostic lumbar punctures were made on the first day of hospitalization and spinal fluids were examined for appearance, protein, sugar and chloride content, cell counts and differentials, and bacteria as revealed by gram stains and cultures. Other procedures included routine blood counts, urinalysis, stool examinations in patients with diarrhea, blood sugar determinations and bacteriological cultures of nasopharynx and blood.

Response to therapy was evaluated by the usual clinical criteria, including duration of fever, improvement of neurologic status, and decrease in peripheral blood leucocyte count with a return of the differential count toward normal and improvement in C.S.F. findings (including pressure, cell count, percentage of polymorphonuclear leucocytes, and glucose and protein content). Outcome of therapy was evaluated according to the patient's courses in the hospital, clinical status at the time of discharge and, whenever possible, a follow-up study one year after discharge.

PROGRESS

Bacterial isolates from CSF by age and sex are shown in Table 1. Most patients were less than one year old and were seriously ill when they arrived at the hospital (Table 2). Eleven of the total of 19 deaths occurred during the first 24 hours of hospitalization. All had high CSF cell counts and, in 9 of the 16 cases of pneumococcal meningitis, there was concomitant septicemia (Table 3). Antibiotic sensitivity studies as determined by tube dilution tests indicated that all isolates of pneumococci were very sensitive *in vitro* to penicillin and ampicillin and all but 2 isolates were sensitive to 1.56 mcg/ml of chloramphenicol. All isolates of *H. influenzae* were sensitive to chloramphenicol and ampicillin and slightly sensitive to kanamycin. All strains were resistant to penicillin G. All isolates of pneumococci and *H. influenzae* were resistant to sulfadiazine. Possible drug reactions or complications of therapy were not noted.

SUMMARY

A study of acute bacterial meningitis in Thai children has been continued. Results to date indicate that the causal agent is *D. pneumoniae* or *H. influenzae* in most cases. While all isolates of *D. pneumoniae* were sensitive *in vitro* to antibiotics in each regimen, the mortality rate was 56.3%. The mortality rate of patients with *H. influenzae* meningitis was 46.2%. There are too few patients in each group to enable meaningful comparisons of therapeutic regimens at this time.

Table 1

Acute Bacterial Meningitis:
Bacteria Isolated from Cerebrospinal Fluids by Age & Sex

Bacteria Isolated	Sex	Total Patients	6 Months	6-12 Months	1-2 Years	2 Years or more	Discharged	Expired
<u>Diplococcus pneumoniae</u>	Male	12	8	3	0	1	6	6
	Female	4	1	3	0	0	1	3
<u>Haemophilus influenzae</u>	Male	5	2	3	0	0	3	2
	Female	8	4	4	0	0	4	4
Others	Male	4	2	0	1	1	3	1
	Female	2	1	0	1	0	2	0
No growth from CSF	Male	8	4	1	0	3	5	3
	Female	1	0	0	1	0	1	0
Total		44	22	14	3	5	25	19

Table 2
Effect of Antimicrobial Therapy on Haemophilus influenzae and Pneumococcal Meningitis

Organism Isolated from Cerebrospinal fluid	Therapy	Clinical Status on Admission to Hospital*	Non Survivors		Survivors			Status on Discharge	
			No. Hospitalized Prior to Death (Average)	Period Hospitalized (Average)	No.	Average Duration of Temp 38°C (Days)	Duration of Hospitalization (Days)	Residual	No Residual
<u>Haemophilus influenzae</u>	Penicillin + Kanamycin	++ +++ ++++ +++++	1	36 hrs	4	3	34	2**	2
	Penicillin + Chloramphenicol	1	3	12 hrs	2	3	20	0	2
	Ampicillin		2	24 hrs	1	0	23	0	1
<u>Diplococcus pneumoniae</u>	Penicillin + Kanamycin		4	32 hrs	1	8	38	0	1
	Penicillin + Chloramphenicol	1	5	24 hrs	4	4	20	0	4
	Ampicillin	2	0	—	2	8	21	0	2

* One or more of the symptoms listed below
 +++++ Coma, shock or semi-coma
 ++++ Convulsions without coma or shock
 ++ Temp 40 C, symptoms for 5 days, complicating disease(s)

** 1 patient with hydrocephalus and
 1 patient with localized neurological damage to hand.

Table 3

Acute Bacterial Meningitis:
Isolation of Organisms from Various Sites

Organism Isolated	Isolation Sites			
	CSF only	CSF and Blood	CSF, Blood & Nasopharynx	Total Patients
<u>Diplococcus pneumoniae</u>	7	7	2	16
<u>Haemophilus influenzae</u>	11	2	0	13
<u>Neisseria meningitidis</u>	0	2	0	2
Group A streptococcus	1	0	0	1
<u>Salmonella tennessee</u>	1	0	0	1
<u>Pseudomonas spp.</u>	2	0	0	2